



US

BO44718/DBO/FtH

HS

**METHOD FOR THE TREATMENT OF OBESITY, OVERWEIGHT AND
FLUCTUATIONS IN BLOOD INSULINE AND/OR GLUCOSE LEVELS**

1. Field of the inventions

5 The present invention relates to method for the prevention or treatment of overweight, obesity or fluctuations in blood insuline and/or glucose levels in mammals, the method comprising the administration to a mammal of an enzyme capable of converting an ingested carbohydrate or a digestion product thereof into one or more absorbable components, wherein the total metabolic caloric value of the absorbable component(s) is less than the metabolic caloric value of the ingested carbohydrate or digestion product thereof. The invention also provides a preparation useful for such treatment.

2. Background of the invention

15 Methods for treatment or prevention of obesity, overweight and fluctuations in blood glucose and/or blood insulin levels known in the art often make use of foodstuffs with reduced caloric value; compositions stimulating metabolism, e.g. by inducing *in vivo* thermogenesis; or compositions providing *in vivo* inhibition of digestive enzyme activity. Many drawbacks are attached to the methods as described above. Especially low caloric diets are particularly undesirable due to the required change in consumption pattern and the adverse taste of many low caloric foodstuffs. The inhibition of digestive enzyme activity has the disadvantage that it often causes flatulence and that its efficacy is seriously influenced by dietary factors.

25 Like obese individuals, also subjects who desire to reduce blood glucose and/or blood insulin fluctuations have to carefully control their diet, e.g. by consuming limited amounts of carbohydrates or by consuming foodstuffs with a low carbohydrate content. The downsides are significant as, for example, low carbohydrate compositions often have a bad taste.

30

In view of the above considerations there is a need for a method which is capable of reducing the metabolic caloric value of ingested carbohydrate containing foodstuffs, but which does not rely on the inhibition of digestive enzyme activity. In addition there is also a need for a method which reduces the impact of ingested carbohydrate containing foodstuffs on blood insuline and/or blood glucose levels without using inhibition of digestive enzyme activity.

US 4,396,602 describes a method of lowering the blood glucose level in mammals. The method comprises administering an enzyme capable of synthesizing sparingly-digestible saccharides from easily-digestible saccharides. The blood glucose level-lowering agent comprises the enzyme capable of synthesizing sparingly-digestible polysaccharides or oligosaccharides from easily-digestible saccharides, such as monosaccharides, oligosaccharides and polysaccharides. Enzymes providing the above effect are dextranucrase and cyclodextrin-synthesizing enzymes.

A major downside of the use of enzymes catalyzing the formation of indigestible polysaccharides and oligosaccharides is that ingestion of such enzymes may cause flatulence. The saccharides formed by the enzymes will not be absorbed by the intestinal cells and be transported to the colon, where these saccharides will be fermented. The fermentation of the oligo- and polysaccharides will result in excessive flatulence. Furthermore, it is questionable whether the conversion to indigestible polysaccharides is truly effective, in particular since a large fraction of the formed indigestible polysaccharides may be converted back to digestible polysaccharides.. It is noted that a significant reduction of carbohydrate absorption will result in a rapid reappearance of appetite, which is likely to result in the early consumption of additional foodstuff.

The known use of enzymes in food preparations has mainly focused on the modification of food ingredients with the purpose of improving bioavailability of active components or of improving digestibility.

The increase of bioavailability of compounds by the addition of enzymes has been described in US 6,099,844, which relates to a method of increasing the yield of

extractable substance from a botanical in the gastrointestinal system of a human being comprising the step of ingesting an enzyme composition comprising a cellulase enzyme and a pectinase enzyme at approximately the same time as a botanical is ingested so that the cellulase and pectinase enzymes degrade the cellulosic and pectin constituents,
5 respectively, contained in the ingested botanical, to obtain an enhanced quantity of extractable substance from the botanical.

US 4,959,212 provides a non-toxic, oxidizing-energizing composition suitable for use as an accelerator of the carbohydrate oxidative degradation metabolic process or the direct
10 oxidation of glucose. Such a composition is said to be effective to reduce the blood glucose concentration in a human body afflicted with diabetes. The composition optionally comprises an enzyme selected from the group consisting of fructose diphosphate aldolase, phosphofructokinase, hexokinase, glucokinase, glucose 6-
15 phosphate dehydrogenase, glucose phosphate isomerase, D-glucose phosphotransferase and mixtures.

3. Summary of the Invention

5 The present invention discloses a novel method for the prevention and/or treatment of obesity, overweight and fluctuations in blood glucose levels and/or blood insulin levels without the above mentioned drawbacks.

10 The current invention provides a method for the prophylactic and curative treatment of overweight, obesity and fluctuations in blood glucose levels and/or blood insulin levels comprising the enteral administration of a preparation containing an effective amount of an enzyme capable of converting an ingested carbohydrate or digestion products thereof into one or more absorbable components, wherein the total metabolic caloric value of the absorbable component(s) is less than the metabolic caloric value of the ingested carbohydrate or digestion product thereof. Thus the present invention effectively provides
15 a method that allows complete digestion of ingested digestible carbohydrates whilst at the same time reducing the actual metabolic caloric value of said ingested carbohydrates.

In the present method digestible carbohydrates are fully digested. Hence, the method according to the present invention will not cause flatulence. The present method does not
20 require a change in diet and will induce a satiety effect.

25 4. Detailed description of the invention

The present invention provides a method of treating or preventing obesity, overweight, fluctuations in blood insulin levels and/or fluctuations in blood glucose levels, said method comprising the enteral administration of an effective amount of a preparation containing an enzyme capable of converting an ingested carbohydrate or digestion
30 product thereof into one or more absorbable components, wherein the total metabolic caloric value of the absorbable component(s) is less than the metabolic caloric value of the ingested carbohydrate or digestion product thereof.

Enzymes

The enzyme used in the method according to the present invention is capable of converting an ingested carbohydrate into one or more absorbable components, wherein the total metabolic caloric value of the absorbable component(s) is below the metabolic caloric value of the ingested carbohydrate. According to a preferred embodiment, the enzyme is selected from the group of isomerases.

A main cause of overweight is the ingestion of vast amounts of glucose monomers or polymers including glucose monomers. According to a preferred embodiment the enzyme is capable of converting glucose into an absorbable component having a decreased metabolic caloric value compared to glucose. Especially advantageous is the conversion of glucose into fructose, thereby additionally providing the benefits of fructose, such as its thermogenic activity and appetite reducing properties. Hence, according to a particularly preferred embodiment, the enzyme is capable of the isomerization of glucose into fructose, i.e. glucose isomerase.

Metabolic caloric value reduction

The term metabolic caloric value as used in the present invention encompasses the caloric value derivable from a carbohydrate by a mammal by complete oxidation of the carbohydrate. The metabolic caloric value of a carbohydrate can be determined on a theoretical basis and by analysis. When the theoretical metabolic caloric value of a carbohydrate is determined, the ATP required for the oxidation of the carbohydrate should be subtracted from the amount of ATP, which the complete oxidation of the carbohydrate would yield in a mammal.

In the case of glucose, absorption of orally ingested glucose in the duodenum requires about 0.5 mol ATP/mol glucose. The blood glucose can subsequently be converted either to fructose 1,6, diphosphate or can enter the gluconeogenesis via uridyldiphosphoglucose (UDPG), requiring 2 mol ATP/mol glucose. Fructose 1,6 diphosphate can subsequently be completely oxidized to form carbon dioxide and water, releasing a total of 38 mol ATP/mol glucose. Net gain of ATP per mol of exogeneous glucose oxidized is therefore 35.5 mol ATP. The metabolic caloric value of glucose is therefor 35.5 mol ATP/mol

glucose.

The metabolic caloric value of fructose is 34.5 mol ATP/mol fructose. Fructose is, similar to glucose, absorbed in the intestine by a process requiring about 0.5 mol ATP/mol fructose. Fructose is generally metabolized essentially in the liver, where the enzyme fructokinase catalyses the phosphorylation of fructose into fructose-1-phosphate, requiring 1 mol ATP per mol of oral fructose. Subsequently, the fructose-1-phosphate is converted to glyceraldehyde phosphate (GAP) and dihydroxyacetone phosphate (DHAP) by the enzyme aldolase B. DHAP can be further degraded to pyruvate and enter the tricarboxylic acid cycle, or can be reconverted into glucose in the process of gluconeogenesis.

The conversion of orally ingested fructose to glycogen, requiring 2 mol ATP per mol of oral fructose (the conversion of glyceraldehyde to glyceraldehyde phosphate (GAP) requires hydrolysis of 1 mol ATP per mol of oral fructose and uridine triphosphate regeneration requires 1 mol ATP). Thus storage of orally ingested fructose as glycogen will require 3.5 mol ATP/mol fructose compared to 2.5 mol ATP/mol glucose.

The energy cost of storing fructose is thus about 9.2 % of the caloric value of fructose, while the energy cost of storing glucose is about 6.5 % of the caloric value of glucose. Consequently, fructose has a substantially lower metabolic caloric value than glucose.

The effect of glucose isomerase on the energy expenditure can be more accurately determined by measurement of energy expenditure by indirect calorimetry according to Scharz et al (Am J Physiol 1992;262(4 Pt 1):E394-401).

Glucose isomerase

Several glucose isomerases with different characteristics are known in the art. According to a preferred embodiment a glucose isomerase is used which shows significant activity at the pH which normally occurs in the duodenum. Preferably the glucose isomerase has a pH optimum for converting glucose to fructose below 8.5, more preferably below 8, even more preferably below 7.5. The optimum is preferably at a pH above 4, even more preferably above 5.

Dosages

In accordance with the present invention the glucose isomerizing enzyme is suitably administered in an amount of between 10 and 100.000 international units (IU) per gram of the dosage. Preferably 0.5 to 1500 international units (IU) enzyme per kg body weight of the mammal are administered to the mammal per dosage. More preferably the dosage includes 1 to 750 IU enzyme per kg body weight, even more preferably 2 to 500 IU enzyme per kg body weight, most preferably 10 to 100 IU enzyme per kg body weight. Preferably the enzyme is glucose isomerase.

The glucose isomerase is preferably administered in a concentrated dosage form. The glucose isomerase can suitably be administered in a preparation preferably comprising between 25 and 10.000 IU glucose isomerase per gram, more preferable between 100 and 5000 IU glucose isomerase per gram, most preferably between 250 and 2500 IU glucose isomerase per gram.

Whenever the tem international unit (IU) is used in the present document this refers to the quantity of enzyme, which transfers 1 micromol glucose per minute to a carbohydrate having a lower caloric value than glucose, at pH 7.5, and 37 °C. For example, 1 IU glucose isomerase refers to the quantity of glucose isomerase, which transfers 1 micromol glucose per minute to fructose at pH 7.5, and 37 °C.

With glucose as a substrate, glucose isomerase activity can be assayed by the measurement of D-fructose produced during the isomerization reaction using the cysteine-carbazole method (CCM) which is based on the reaction of ketosugars with carbazole in acids to yield a purple product (Dische and Borenfreund, J. Biol. Chem. 192 (1951) 583).

Whenever the term dose or dosage is used within this disclosure, any dosage form is encompassed which can be administered enterally (e.g. orally), within a fairly narrow time span. Whenever reference is made to a certain quantity that is administered per dose or dosage, said quantity is preferably administered within one hour, more preferably within 15 minutes, even more preferably within 5 minutes.

Preparation

The term preparation within the spirit of the present invention refers to nutritional as well as pharmaceutical compositions. Pharmaceutical compositions may suitably include a pharmaceutically acceptable carrier. Pharmaceutical acceptable carriers are well known and described in the art. The preparation used in the present method can be applied in any suitable form, such as meals, bars, pills, capsules, gels, biscuits, drinks etc. According to a preferred embodiment the preparation is administered in a solid or semisolid dosage form, more preferably in the form of a pill, which term includes capsules, tablets, microparticles and microspheres.

The aforementioned single solid or semisolid dosage form preferably has a weight between 0.1 and 30 grams, more preferably between 0.2 and 10 gram. When a pill is used to provide the enzyme, the pill preferably has a weight between 0.2 and 4 grams, even more preferably between 0.5 and 3 grams. A dosage can include one or more pills, however, preferably the dosage consists of 1 to 3 pills.

Many enzymes will not survive the acidic environment of the stomach. The enzyme used in the present method is preferably administered in an pill that is coated with a substance that can withstand the enteric environment (an enteric coating) or in another form that prevents the decrease of enzyme activity, e.g. by co-administering a buffer and/or by co-administering inhibitors of intestinal proteolytic enzymes. Alternatively or additionally, enzymes may be used which have reduced sensitivity to proteolytic breakdown or which are not or only partially affected by an acidic environment.

According to a particularly preferred embodiment the enzyme is administered in a solid or semi-solid dosage form with a coating that prevents the reduction of activity of the enzymes by stomach acid and/or stomach proteases. A delayed, post-gastric, release of the active enzymes in the small intestine (duodenum, ileum, jejunum) can be achieved by encasing the enzymes. One class of acid-resistant agents suitable for this purpose is that disclosed in Eury et al., U.S. Pat. No. 5,316,774. Effective enteric materials include polyacids having a pK_a of from about 3 to 5. Examples of such materials are fatty acid mixtures, methacrylic acid polymers and copolymers, ethyl cellulose, and cellulose

acetate phthalates. Specific examples are methacrylic acid copolymers sold under the name EUDRAGIT.RTM., available from Rohm Tech, Inc., Maiden, Mass., USA; and the cellulose acetate phthalate latex AQUATERIC.RTM., available from FMC Corporation, New York, N.Y., USA, and similar products available from Eastman-Kodak Co.,
5 Rochester, N.Y., USA.

Thermogenic effect of fructose

According to a preferred embodiment of the present method fructose is generated from ingested glucose. Fructose has been shown to provide an increased thermogenic effect compared to glucose. It is the inventors belief that an additional energy expenditure is required during the metabolisms of fructose, even further decreasing the metabolic caloric value of fructose. Ingestion of an enzyme capable of converting glucose into fructose, e.g. glucose isomerase, will therefore induce a thermogenic effect. This thermogenic effect contributes to the prevention or treatment of obesity or overweight. (Schwarz et al; Thermogenesis in obese women: effect of fructose vs. glucose added to a meal. Am J Physiol 1992;262(4 Pt 1):E394-401.)
10
15

Appetite reducing effect of fructose

20 Furthermore, fructose ingestion is suggested to decrease food intake. Several mechanisms have been suggested to cause this appetite suppressing effect, however, the mechanism has not been elucidated. The suggested appetite reducing effect induced by fructose might be caused by the effect fructose has on gastric emptying. Fructose empties in a rapid, exponential fashion, while glucose empties in a more slowly, linear fashion.
25 However, a more likely explanation for the appetite reducing effect of fructose can be found in the reduced fluctuation in plasma insulin levels and/or plasma glucose levels. Fructose ingestion leads to lower values of insulin in comparison to glucose ingestion. High insulin concentrations have been related to hunger feelings. Furthermore, there is evidence that glucose and fructose have a different impact on hepatic metabolism, which
30 metabolism is believed to influence food intake. In animal studies it has been shown that jugular infusion of fructose, as opposed to glucose, decreases food intake when given

before eating.

Ingestion of an enzyme converting glucose into fructose, e.g. glucose isomerase, will therefore reduce appetite and prevent hunger. The reduction of appetite is a highly desired impact for a preparation that is used in a method for the prevention and/or treatment of obesity or overweight. An enzyme capable of converting glucose into fructose is therefore especially useful in the method for the prevention and/or treatment of overweight and obesity.

Under normal physiological circumstances, ingested digestible di-, tri-, or polysaccharides are converted into monosaccharides in the acidic environments and/or by the carbohydrase activity in the mammalian intestinal tract. The monosaccharides are subsequently absorbed by the cells in the duodenum. Whenever reference is made in this document to ingested carbohydrate or a digestion product thereof, monosaccharides as well as digestible di-, tri-, oligo- or polysaccharides which can be converted into monosaccharides in the gastro-intestinal tract are meant.

Preferably, the absorbable carbohydrate formed by the enzyme used in the method according to the invention has a molecular weight between 75% and 125 % of the molecular weight of the substrate, i.e. the ingested carbohydrate or digestion product thereof, preferably between 90% and 110%, even more preferably between 95% and 105%, especially between 99% and 101%. According to an especially preferred embodiment, the ingested carbohydrate or digestion product thereof is glucose or a di-, tri-, oligo- or polysaccharide containing glucose monose units and the absorbable carbohydrate is fructose.

In compositions meant for weight control, treatment or prevention of obesity or overweight often glucose has been (partially) replaced by fructose because of the above reasons. Although such diets provide at least part of the desired effects of fructose, still a vast amount of carbohydrates are consumed in such diets. Exclusion of "all" glucose comprising di-, tri-, and polysaccharides from foodstuff is impossible, in view of technical and commercial considerations. It is therefor desirable to accomplish the above advantageous effects of fructose, without the need of ingesting relatively large quantities

of fructose. As explained herein before, this can be achieved by the ingestion of an enzyme capable of converting glucose to a monosaccharide of lower metabolic caloric value, e.g. fructose, such an enzyme preferably being a glucose isomerase.

5 *Prevention and treatment of blood glucose fluctuations*

The glycemic index is a measure for the effect of ingested foodstuff on blood glucose levels. The index gives a relative value for the blood sugar increase following the ingestion of the foodstuff. Fructose has a lower glycemic index (GI) value, compared to glucose (GI value glucose = 100; GI value fructose = 20). Additionally, fructose is first
10 absorbed in the small intestine, then transported to the liver for conversion to glucose, its initial uptake is insulin independent.

Diabetics must manage their diet to maintain a normal blood glucose level: any increase in blood glucose will trigger an insulinemic response, creating an imbalance. This could lead to a serious insulin reaction or coma. Fructose, unlike glucose, does not cause a high
15 initial glucose spike.

The use of enzymes capable of converting ingested glucose monosaccharides into fructose prevents abnormal insulin levels, reduces the insulinemic response of ingested glucose monosaccharide and provides a decreased fluctuation in blood glucose levels, all of which are highly desirable for subjects suffering from diabetes and associated diseases.

20 Thus the present method may advantageously be used in the treatment or prevention of fluctuations in blood glucose levels and related disorders such as abnormal insulin levels, major fluctuations in blood insulin levels, insulinemic response after ingestion of foodstuff.

25 *Cofactors/inhibitors*

According to a preferred embodiment, the preparation further contains cofactors, e.g. minerals, that increase the activity of the enzyme. When glucose isomerase is used, preferably magnesium and/or cobalt is coadministered. Magnesium can be included in the composition containing glucose isomerase in an amount between 10 mg and 5 g per
30 dosage, more preferably between 30 mg and 1 g, even more preferably between 40 mg and 450 mg.

Combinations

In order to further improve the present method, the enzyme may be coadministered with components capable of decreasing the absorption or digestion of ingested carbohydrates or digestion products thereof, e.g. carbohydrase inhibitors. Co-administration of such ingredients will increase the retention time of ingested and (partially) digested carbohydrate material in the duodenum, thereby increasing the amount of absorbable monosaccharide formed from the ingested carbohydrate per unit active enzyme. Preferred carbohydrate absorption inhibitors are gymnemic acid (e.g. obtainable from *gymnema*) or soluble indigestible fibers such as glucomannan and locust bean gum. Preferred carbohydrase inhibitors include plant derived polyphenols, selected from the group of catechins or derivatives thereof, anthocyanidins, proanthocyanidins, procyanidins and cyanidins, which are exemplary and preferably obtained from green tea (*Camellia sinensis*) or grape (*Vitis vinifera*). The above components may be coadministered with the present enzyme in an amount of 0.001 to 1000 mg/IU of the enzyme, more preferably 0.01 to 100 mg/IU of the present enzyme.

Application

The enzyme is preferably administered to mammals having a body weight above 25 kg, more preferably to humans. Furthermore, the preparation can be advantageously used in the manufacture of a medicament for use in a method for the treatment and prevention of obesity or overweight, the method comprising the administration of an effective amount of glucose isomerase to a human.

A further objective of the present invention is to provide a cosmetic method for reducing or preventing the formation of body fat or keeping a lean body, comprising administering a therapeutically effective amount of a preparation comprising an enzyme capable of converting an ingested carbohydrate or digestion product thereof into one or more absorbable components, wherein the total metabolic caloric value of the absorbable component(s) is below the metabolic caloric value of the ingested carbohydrate or digestion product thereof.

The enzymes are preferably administered between 60 minutes before and 60 minutes

after the ingestion of a significant amount of carbohydrates, e.g. at least 5 grams of carbohydrates. According to a further preferred embodiment, the enzyme is ingested prior to, during or shortly after a meal.

The enzymes are preferably ingested in the form of a pharmaceutical preparation or as a

5 nutritional supplement.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
22

5. Examples

Example 1: Pharmaceutical composition

5

A tablet having an outside coating consisting of EUDRAGIT.RTM containing:
1 gram glucose isomerase (glucose isomerase 350 IGIU/gram, Sweetzyme T, Novozymes A/S, Denmark) and 150 mg magnesium chloride

Example 2: Nutritional supplement

A nutritional supplement in the form of a gelatin capsule advertised to decrease the caloric value of ingested foodstuff and/or decrease blood glucose fluctuations comprising:

750 mg glucose isomerase (1500 IGIU/ml glucose isomerase (G-zyme, G993, obtained from Enzyme Bio-Systems, Beloit, USA)) and
250 mg Gymnema Sylvestre extract (comprising 15wt% gymnemic acid)

Example 3: Fructose formation under sub-intestinal conditions

To test the fructose forming properties of glucose isomerase under intestinal conditions mixtures of: 5 ml starch solution (5 ml 7.5 g Pacelli potato starch /100 ml 50 mM phosphate buffer; Paselli WA4 potato starch, AVEBE, Foxhol, The Netherlands), amylase (1 ml A6211, obtained from Sigma Chemie, Zwijndrecht) and brush border enzymes (0.2 ml, scraping of the inner wall of piglet small intestinal wall) were prepared. The mixtures were adjusted to pH 6.5 using 2 ml 50 mM phosphate buffers, which mimics the pH in the human intestine (pH 6-7.5). A mixture with and without 0.18 ml glucose isomerase (G-zyme, G993, obtained from Enzyme Bio-Systems, Beloit, USA) was incubated and the concentration of glucose and fructose was measured over time.

Table 1 gives the concentration glucose and fructose in the mixtures with and without glucose isomerase in time.

TABLE 1

Time (hours)	Without glucose isomerase	With glucose isomerase			
		Glucose concentration (g/l)	Fructose concentration (g/l)	Glucose + Fructose concentration (g/l)	Conversion rate glucose to fructose (%)
0	2.6	1.8	0	1.8	0
0.5	6.3	4.1	1.5	5.6	26
1	10.6	6.8	3.0	9.8	31
3	13.8	9.2	4.6	13.8	33
4	15.5	10	5.6	15.5	36
6	15.6	12.3	7.0	19.4	36

From this study it can be concluded that fructose formation from glucose takes place under conditions as present in the small intestine.

Example 4: Fructose formation under intestinal conditions using pancreatine

To test the fructose forming properties of glucose isomerase under intestinal conditions, in the presents of pancreatine (including pancreas proteases), a mixture of 5 ml starch solution (7.5 g Pacelli potato starch /100 ml 50 mM phosphate buffer; Paselli WA4 potato starch, AVEBE, Foxhol, The Netherlands), 1.75 gram pancreatine (P1750, Sigma Chemie, Zwijndrecht), 475 mg cow bile and 0.15 ml brush border enzymes (scraping of the inner wall of piglet small intestinal wall) was prepared. The mixtures were adjusted to pH 6.5 using 2 ml 50 mM phosphate buffers, which mimics the pH in the human intestine (pH 6-7.5). A mixture with and without 0.2 ml glucose isomerase (G-zyne, G993, obtained from Enzyme Bio-Systems, Beloit USA) was incubated and the concentration of glucose and fructose was measured over time.

Table 2 gives the concentration glucose and fructose in the mixtures with and without glucose isomerase in time.

5

TABLE 2

Time (hours)	Without glucose isomerase	With glucose isomerase			
		Glucose concentration (g/l)	Fructose concentration (g/l)	Glucose + Fructose concentration (g/l)	Conversion rate glucose to fructose (%)
0	1.5	1.5	0	1.5	0
1	3.7	3.1	0.7	3.7	19
3	5.9	4.5	1.2	5.7	21
7	9.3	7.1	3.4	10.5	32

10

From this study it can be concluded that fructose formation from glucose takes place in the presents of pancreatic proteases.